

## ***Biosynthesis of the respiratory toxin bongkreikic acid in the pathogenic bacterium Burkholderia gladioli.***

Moebius N, Ross C, Scherlach K, Rohm B, Roth M, Hertweck C (2012) *Biosynthesis of the respiratory toxin bongkreikic acid in the pathogenic bacterium Burkholderia gladioli*. *Chem Biol* 19(9), 1164-1174.

[Details](#)



### **Abstract**

Bongkreikic acid (BA), an infamous respiratory toxin of the pathogenic bacterium *Burkholderia gladioli*, causes lethal intoxications when tempe bongkrek is produced with contaminated *Rhizopus oligosporus* cultures. Genome sequencing of *B. gladioli* pathovar *cocovenenans* unveiled the genetic basis for BA biosynthesis, and pointed to a homologous *bon* gene cluster in a *B. gladioli* strain from an infected rice plant. For functional genetics in *B. gladioli*  $\lambda$  Red recombination was established. Dissection of the modular type I polyketide synthase (a trans-AT PKS) provided insights into complex polyketide assembly. Isoprenoid-like  $\beta$ -branching events and a six-electron oxidation of a methyl group to a carboxylic acid give rise to the unique branched tricarboxylic fatty acid. The role of the cytochrome P450 monooxygenase, *BonL*, was proven by structural elucidation of deoxybongkreikic acid from a mutant.

## Beteiligte Forschungseinheiten

[Biomolekulare Chemie Christian Hertweck](#) [Mehr erfahren](#)

[Biotechnikum Miriam Agler-Rosenbaum](#) [Mehr erfahren](#)

## Leibniz-HKI-Autor\*innen



Christian Hertweck

[Details](#)



**Martin Roth**

[Details](#)



**Kirstin Scherlach**

[Details](#)

## Identifizier

**doi:** 10.1016/j.chembiol.2012.07.022

**PMID:** 22999884