

Biosynthesis and function of gliotoxin in *Aspergillus fumigatus*.

Scharf DH, Heinekamp T, Remme N, Hortschansky P, Brakhage AA, Hertweck C (2012) Biosynthesis and function of gliotoxin in *Aspergillus fumigatus*. *Appl Microbiol Biotechnol* 93(2), 467-472.

[Details](#)



Abstract

Gliotoxin (GT) is the prototype of the epidithiodioxopiperazine (ETP)-type fungal toxins. GT plays a critical role in the pathobiology of *Aspergillus fumigatus*. It modulates the immune response and induces apoptosis in different cell types. The toxicity has been attributed to the unusual intramolecular disulfide bridge, which is the functional motif of all ETPs. Because of the extraordinary structure and activity of GT, this fungal metabolite has been the subject of many investigations. The biosynthesis of GT involves unprecedented reactions catalysed by recently discovered enzymes. Here, we summarize the recent progress in elucidating the GT biosynthetic pathway and its role in virulence.

Beteiligte Forschungseinheiten

[Biomolekulare Chemie](#) [Christian Hertweck](#) [Mehr erfahren](#)

[Molekulare und Angewandte Mikrobiologie](#) [Axel Brakhage](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Axel A. Brakhage

[Details](#)



Thorsten Heinekamp

[Details](#)



Christian Hertweck

[Details](#)



Peter Hortschansky

[Details](#)



Daniel H. Scharf

[Details](#)

Identifier

doi: 10.1007/s00253-011-3689-1

PMID: 22094977