

A dual expression platform to optimize the soluble production of heterologous proteins in the periplasm of *Escherichia coli*.

Kraft M, Knüpfer U, Wenderoth R, Kacholdt A, Pietschmann P, Hock B, Horn U (2007) A dual expression platform to optimize the soluble production of heterologous proteins in the periplasm of *Escherichia coli*. *Appl Microbiol Biotechnol* 76(6), 1413-1422.

[Details](#)



Abstract

The functional analysis of individual proteins or of multiprotein complexes - since the completion of several genome sequencing projects - is in focus of current scientific work. Many heterologous proteins contain disulfide-bonds, required for their correct folding and activity, and therefore, need to be transported to the periplasm. The production of soluble and functional protein in the periplasm often needs target-specific regulatory genetic elements, leader peptides, and folding regimes. Usually, the optimization of periplasmic expression is a step-wise and time-consuming procedure. To overcome this problem we developed a dual expression system, containing a degP-promoter-based reporter system and a highly versatile plasmid set. This combines the differential protein expression with the selection of a target-specific expression plasmid. For the validation of this expression tool, two different molecular formats of a recombinant antibody directed to the human epidermal growth factor receptor and human 11beta-hydroxysteroid dehydrogenase type 2 (11beta-HSD2) were used. By application of this expression system we demonstrated that the

amount of functional protein is inversely proportional to the on-line luciferase signal. We showed that this technology offers a simple tool to evaluate and improve the yield of functionally expressed proteins in the periplasm, which depends on the used regulatory elements and folding strategies.

Beteiligte Forschungseinheiten

[Biotechnikum Miriam Agler-Rosenbaum](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Uwe Horn

[Details](#)



Uwe Knüpfer

[Details](#)



Mario Kraft

[Details](#)

Identifier

doi: 10.1007/s00253-007-1121-7

PMID: 17684739