

Human Invariant Natural Killer T cells possess immune-modulating functions during *Aspergillus* infection.

Beitzen-Heineke A, Bouzani M, Schmitt AL, Kurzai O, Hünninger K, Einsele H, Loeffler J (2016) Human Invariant Natural Killer T cells possess immune-modulating functions during *Aspergillus* infection. *Med Mycol* 54(2), 169-176.

[Details](#)



Abstract

Aspergillus fumigatus is the most common cause for invasive fungal infections, a disease associated with high mortality in immune-compromised patients. CD1d-restricted invariant natural killer T (iNKT) cells compose a small subset of T cells known to impact the immune response toward various infectious pathogens. To investigate the role of human iNKT cells during *A. fumigatus* infection, we studied their activation as determined by CD69 expression and cytokine production in response to distinct fungal morphotypes in the presence of different CD1d(+) antigen presenting cells using flow cytometry and multiplex enzyme-linked immunosorbent assay (ELISA). Among CD1d(+) subpopulations, CD1d(+)CD1c(+) mDCs showed the highest potential to activate iNKT cells on a per cell basis. The presence of *A. fumigatus* decreased this effect of CD1d(+)CD1c(+) mDCs on iNKT cells and led to reduced secretion of TNF- α , G-CSF and RANTES. Production of other Th1 and Th2 cytokines was not affected by the fungus, suggesting an immune-modulating function for human iNKT cells during *A. fumigatus* infection.

Beteiligte Forschungseinheiten

[Fungal Septomics](#) [Oliver Kurzai](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Kerstin Hänniger

[Details](#)



Oliver Kurzai

[Details](#)

Identifier

doi: myv074

PMID: 26483428

