

18 F-Fluoride positron emission tomography/computed tomography for noninvasive in vivo quantification of pathophysiological bone metabolism in experimental murine arthritis.

Irmeler IM, Gebhardt P, Hoffmann B, Opfermann T, Figge MT, Saluz HP, Kamradt T (2014) 18 F-Fluoride positron emission tomography/computed tomography for noninvasive in vivo quantification of pathophysiological bone metabolism in experimental murine arthritis. *Arthritis Res Ther* 16(4), R155.

[Details](#)



Abstract

Evaluation of disease severity in experimental models of rheumatoid arthritis is inevitably associated with assessment of structural bone damage. A noninvasive imaging technology allowing objective quantification of pathophysiological alterations of bone structure in rodents could substantially extend the methods used to date in preclinical arthritis research for staging of autoimmune disease severity or efficacy of therapeutical intervention. Sodium 18 F-fluoride (18 F-NaF) is a bone-seeking tracer well-suited for molecular imaging. Therefore, we systematically examined the use of 18 F-NaF positron emission tomography/computed tomography (PET/CT) in mice with glucose-6-phosphate isomerase (G6PI)-induced arthritis for quantification of

pathological bone metabolism.

Beteiligte Forschungseinheiten

[Zell- und Molekularbiologie](#) [Mehr erfahren](#)

[Angewandte Systembiologie](#) [Marc Thilo Figge](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Marc Thilo Figge

[Details](#)



Bianca Hoffmann

[Details](#)



Ingo Irmeler

[Details](#)



Hans Peter Saluz

[Details](#)

Identifier

doi: 10.1186/ar4670

PMID: 25053370