

Genetic engineering activates biosynthesis of aromatic fumaric acid amides in the human pathogen *Aspergillus fumigatus*.

Kalb D, Heinekamp T, Lackner G, Scharf DH, Dahse HM, Brakhage AA, Hoffmeister D (2015) Genetic engineering activates biosynthesis of aromatic fumaric acid amides in the human pathogen *Aspergillus fumigatus*. *Appl Environ Microbiol* 81, 1594-1600.

[Details](#)



Abstract

The *Aspergillus fumigatus* nonribosomal peptide synthetase FtpA is among the few of this species whose natural product has remained unknown. Both FtpA adenylation domains were characterized in vitro. Fumaric acid was identified as preferred substrate of the first and both L-tyrosine and L-phenylalanine as preferred substrates of the second adenylation domain. Genetically engineered *A. fumigatus* strains expressed either ftpA or the regulator gene ftpR, encoded in the same cluster of genes, under the control of the doxycycline-inducible tet-on cassette. These strains produced fumaryl-L-tyrosine and fumaryl-L-phenylalanine which were identified by liquid chromatography and high-resolution mass spectrometry. Modeling of the first adenylation domain in silico provided insight into the structural requirements to bind fumaric acid as peptide synthetase substrate. This work adds aromatic fumaric acid amides to the secondary metabolome of the important human pathogen *A. fumigatus* which was previously not known as a producer of these compounds.

Beteiligte Forschungseinheiten

[Infektionsbiologie Peter F. Zipfel](#) [Mehr erfahren](#)

[Molekulare und Angewandte Mikrobiologie Axel Brakhage](#) [Mehr erfahren](#)

[Pharmazeutische Mikrobiologie Dirk Hoffmeister](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Axel A. Brakhage

[Details](#)



Hans-Martin Dahse

[Details](#)



Thorsten Heinekamp

[Details](#)



Dirk Hoffmeister

[Details](#)



Daniel Kalb

[Details](#)



Gerald Lackner

[Details](#)



Daniel H. Scharf

[Details](#)

Identifier

doi: pii: AEM.03268-14

PMID: 25527545