

Clinical-scale isolation of the total *Aspergillus fumigatus*-reactive T-helper cell repertoire for adoptive transfer.

Bacher P, Jochheim-Richter A, Mockel-Tenbrink N, Kniemeyer O, Wingenfeld E, Alex R, Ortigao A, Karpova D, Lehrnbecher T, Ullmann AJ, Hamprecht A, Cornely O, Brakhage AA, Assenmacher M, Bonig H, Scheffold A (2015) Clinical-scale isolation of the total *Aspergillus fumigatus*-reactive T-helper cell repertoire for adoptive transfer. *Cytotherapy* 17(10), 1396-1405.

[Details](#)



Abstract

Evidence of the criticality of the adaptive immune response for controlling invasive aspergillosis has been provided. This observation is supported by the fact that invasive aspergillosis, a grave complication of allogeneic stem cell transplantation, occurs long after myeloid reconstitution in patients with low T-cell engraftment and/or on immunosuppressants. Adoptive T-cell transfer might be beneficial, but idiosyncrasies of *Aspergillus fumigatus* and the anti-*Aspergillus* immune response render established selection technologies ineffective.

Beteiligte Forschungseinheiten

[Molekulare und Angewandte Mikrobiologie Axel Brakhage](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Axel A. Brakhage

[Details](#)



Olaf Kniemeyer

[**Details**](#)

Identifier

doi: S1465-3249(15)00926-3

PMID: 26188965