

Comparative genomics of serial *Candida glabrata* isolates and the rapid acquisition of echinocandin resistance during therapy.

Barber AE*, Weber M, Kaerger K, Linde J, Gölz H, Duerschmied D, Markert A, Guthke R, Walther G, Kurzai O (2019) Comparative genomics of serial *Candida glabrata* isolates and the rapid acquisition of echinocandin resistance during therapy. *Antimicrob Agents Chemother* 63(2), e01628-18.

[Details](#)

*equal contribution



Abstract

The opportunistic pathogen *Candida glabrata* shows a concerning increase in drug resistance. Here we present the analysis of two serial bloodstream isolates, taken 12 days apart. Both isolates show pan-azole resistance and echinocandin resistance was acquired during the sampling interval. Genome sequencing identified 9 nonsynonymous SNVs between the strains, including a S663P substitution in FKS2 and previously undescribed SNVs in MDE1 and FPR1, offering insight

into how *C. glabrata* acquires drug resistance and adapts to a human host.

Beteiligte Forschungseinheiten

[Fungal Septomics Oliver Kurzai](#) [Mehr erfahren](#)

[Microbiome Dynamics Gianni Panagiotou](#) [Mehr erfahren](#)

[Nationales Referenzzentrum für Invasive Pilzinfektionen Oliver Kurzai](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Amelia Barber

[Details](#)



Reinhard Guthke

[Details](#)



Kerstin Kaerger

[Details](#)



Oliver Kurzai

[Details](#)



Jörg Linde

[Details](#)



Grit Walther

[Details](#)



Michael Weber

[Details](#)

Themenfelder

[Systembiologie von Erkrankungen](#)

Identifizier

doi: 10.1128/AAC.01628-18

PMID: 30478162