

Bioactivity and mode of action of bacterial tetramic acids.

Klapper M, Paschold A, Zhang S, Weigel C, Dahse HM, Götze S, Pace S, König S, Rao Z, Reimer L, Werz O, Stallforth P (2019) Bioactivity and mode of action of bacterial tetramic acids. *ACS Chem Biol* 14(8), 1693-1697.

[Details](#)



Abstract

Microbially produced 3-acyltetramic acids display a diverse range of biological activities. The pyreudiones are new members of this class that were isolated from bacteria of the genus *Pseudomonas*. Here, we performed a structure-activity relationship study and determined their mode of action. An efficient biomimetic synthesis was developed to synthesize pyreudione A. Pyreudiones and synthetic analogs thereof were tested for their amoebicidal, antibacterial, antiproliferative, and cytotoxic activities. The length of the alkyl side chain and the nature of the amino acid residues within the tetramic acid moiety strongly affected activity, in particular against mycobacteria. The mode of action was shown to correlate with the ability of pyreudiones to act as protonophores. Removal of the acidic proton by methylation of pyreudione A resulted in a loss of bioactivity.

Beteiligte Forschungseinheiten

[Infektionsbiologie Peter F. Zipfel](#) [Mehr erfahren](#)

[Molekulare und Angewandte Mikrobiologie Axel Brakhage](#) [Mehr erfahren](#)

[Paläobiotechnologie Pierre Stallforth](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Hans-Martin Dahse

[Details](#)



Sebastian Götze

[Details](#)



Martin Klapper

[Details](#)



André Paschold

[Details](#)



Lisa Reimer

[Details](#)



Pierre Stallforth

[Details](#)



Christiane Weigel

[Details](#)

Identifier

doi: 10.1021/acschembio.9b00388

PMID: 31294961