

Pathogenesis of *Candida albicans* infections in the alternative chorio-allantoic membrane chicken embryo model resembles systemic murine infections.

Jacobsen ID, Grosse K, Berndt A, Hube B (2011) Pathogenesis of *Candida albicans* infections in the alternative chorio-allantoic membrane chicken embryo model resembles systemic murine infections. *PLOS One* 6(5), e19741-e19741.

[Details](#)



Abstract

Alternative models of microbial infections are increasingly used to screen virulence determinants of pathogens. In this study, we investigated the pathogenesis of *Candida albicans* and *C. glabrata* infections in chicken embryos infected via the chorio-allantoic membrane (CAM) and analyzed the virulence of deletion mutants. The developing immune system of the host significantly influenced susceptibility: With increasing age, embryos became more resistant and mounted a more balanced immune response, characterized by lower induction of proinflammatory cytokines and increased transcription of regulatory cytokines, suggesting that immunopathology contributes to pathogenesis. While many aspects of the chicken embryo response resembled murine infections, we also observed significant differences: In contrast to systemic infections in mice, IL-10 had a beneficial effect in chicken embryos. IL-22 and IL-17A were only upregulated after the peak mortality in the chicken embryo model occurred; thus, the role of the Th17 response in this model remains unclear. Abscess formation occurs frequently in murine models, whereas the avian

response was dominated by granuloma formation. Pathogenicity of the majority of 15 tested *C. albicans* deletion strains was comparable to the virulence in mouse models and reduced virulence was associated with significantly lower transcription of proinflammatory cytokines. However, fungal burden did not correlate with virulence and for few mutants like *bcr1*Δ and *tec1*Δ different outcomes in survival compared to murine infections were observed. *C. albicans* strains locked in the yeast stage disseminated significantly more often from the CAM into the embryo, supporting the hypothesis that the yeast morphology is responsible for dissemination in systemic infections. These data suggest that the pathogenesis of *C. albicans* infections in the chicken embryo model resembles systemic murine infections but also differs in some aspects. Despite its limitations, it presents a useful alternative tool to pre-screen *C. albicans* strains to select strains for subsequent testing in murine models.

Beteiligte Forschungseinheiten

[Mikrobielle Pathogenitätsmechanismen](#) [Bernhard Hube](#) [Mehr erfahren](#)

[Mikrobielle Immunologie](#) [Ilse Jacobsen](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Katharina Große

[Details](#)



Bernhard Hube

[Details](#)



Ilse Denise Jacobsen

[Details](#)

Identifier

doi: 10.1371/journal.pone.0019741

PMID: 21603634