

The yeast *Candida albicans* evades human complement attack by secretion of aspartic proteases.

Gropp K, Schild L, Schindler S, Hube B, Zipfel PF, Skerka C (2009) The yeast *Candida albicans* evades human complement attack by secretion of aspartic proteases. *Mol Immunol* 47(2-3), 465-475.

[Details](#)



Abstract

Candida albicans, which represents one of the most important human pathogenic yeasts, is directly attacked by the host innate immune system upon infection. However this pathogen has developed multiple strategies to escape host immune defense. Here, we show that *C. albicans* secreted proteases interfere and inactivate host innate immune effector components, such as complement proteins. Secreted aspartic proteases (Saps) in the culture supernatant of *C. albicans* cells and also recombinant Sap1, Sap2 and Sap3 degrade host complement components C3b, C4b and C5 and also inhibit terminal complement complex (TCC) formation. This proteolytic activity is specific to the three recombinant and wild type Sap proteins. The triple knock out *C. albicans* strain Delta sap1-3 and also the non-pathogenic yeast *S. cerevisiae* lack such degrading activities. The complement inhibitory role of Sap1, Sap2 and Sap3 was confirmed in hemolysis assays with rabbit erythrocytes and normal human plasma. Secretion of complement degrading proteases provides a highly efficient complement defense response of this human pathogenic yeast that acts after the immediate acquisition of host complement regulators to the cell surface.

Beteiligte Forschungseinheiten

[Infektionsbiologie Peter F. Zipfel](#) [Mehr erfahren](#)

[Mikrobielle Pathogenitätsmechanismen Bernhard Hube](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Bernhard Hube

[Details](#)



Lydia Kasper

[Details](#)



Christine Skerka

[Details](#)



Peter F. Zipfel

[Details](#)

Identifier

doi: 10.1016/j.molimm.2009.08.019

PMID: 19880183