

Different stimuli for inference of gene regulatory network in rheumatoid arthritis

Kupfer P, Vlaic S, Huber R, Kinnne RW, Guthke R (2013) Different stimuli for inference of gene regulatory network in rheumatoid arthritis In: Pedro Fernandes, Jordi Solé-Casals, Ana Fred and Hugo Gamboa (eds.) Bioinformatics 2013 - 4th Int. Conf. on Bioinformatics Models, Methods and Algorithms International Conference on Bioinformatics Models, Methods and Algorithms, Barcelona /Spain, 02/11/2013-02/14/2013, pp. 282-287.SCITEPRESS – Science and Technology Publications, Lissabon /Portugal. ISBN: 978-989-8565358.

[Details](#)




Abstract

Since genetic and epigenetic factors are known to be involved in the pathogenesis of rheumatoid arthritis the search for key players in this disease is one of the most important challenges. For this purpose gene regulatory networks are one possibility to reveal underlying interactions for different stimuli. In this study we analyzed the cellular response of synovial fibroblasts to 4 different stimuli. We inferred a gene regulatory network that is able to explain the observed data for stimulation by TNF- α , TGF- β 1, IL-1 and PDGF-D simultaneously.

Beteiligte Forschungseinheiten

[Microbiome Dynamics](#) [Gianni Panagiotou](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Reinhard Guthke

[Details](#)



Peter Kupfer

[Details](#)



Sebastian Vlaic

[Details](#)

Themenfelder

[Netzwerke der Wirt-Pilz-Pathogen Interaktion](#)