

Characterization of *Burkholderia rhizoxinica* and *B. endofungorum* isolated from clinical specimens.

Gee JE, Glass MB, Lackner G, Helsel LO, Daneshvar M, Hollis DG, Jordan J, Morey R, Steigerwalt A, Hertweck C (2011) Characterization of *Burkholderia rhizoxinica* and *B. endofungorum* isolated from clinical specimens. *PLOS One* 6(1), e15731.

[Details](#)



Abstract

Eight isolates submitted to CDC from 1989 to 2006 from clinical specimens were initially identified as members of the genus *Burkholderia* based on preliminary cellular fatty acid analysis and/or 16S rRNA gene sequencing. With the recent descriptions of the new species *B. rhizoxinica* and *B. endofungorum*, which are considered endosymbiotic bacteria in *Rhizopus microsporus* fungi, we now identify seven of these clinical isolates as *B. rhizoxinica* and one as *B. endofungorum* based on biochemical testing, 16S rRNA, and DNA-DNA hybridization results. We also further characterize these isolates by assessing toxin production and/or by multiple locus sequence typing.

Beteiligte Forschungseinheiten

[Biomolekulare Chemie](#) [Christian Hertweck](#) [Mehr erfahren](#)

Leibniz-HKI-Autor*innen



Christian Hertweck

[Details](#)



Gerald Lackner

[Details](#)

Identifier

doi: 10.1371/journal.pone.0015731

PMID: 21267449