

# ***Sporothrix humicola* (Ascomycota: Ophiostomatales) - A soil-borne fungus with pathogenic potential in the eastern quoll (*Dasyurus viverrinus*).**

Nessler A, Schauerte N, Geiger C, Kaerger K, Walther G, Kurzai O, Eisenberg T (2019) *Sporothrix humicola* (Ascomycota: Ophiostomatales) - A soil-borne fungus with pathogenic potential in the eastern quoll (*Dasyurus viverrinus*). *Med Mycol Case Rep* 25, 39-44.

[Details](#)



## **Abstract**


The genus *Sporothrix* contains both species pathogenic to humans and animals as well as environmental fungi. *S. humicola*, a member of the latter *S. pallida* clade, has previously been reported only from soil. We have isolated this dimorphic fungus from multiple cutaneous lesions in two endangered marsupials native to Tasmania. Clinical appearance resembled cutaneous sporotrichosis, highlighting the principle pathogenic potential. Identification was based on partial ITS,  $\beta$ -tubulin and calmodulin gene sequencing.

## Involved units

[Fungal Septomycosis Oliver Kurzai](#) [Read more](#)

[National Reference Center for Invasive Fungal Infections Oliver Kurzai](#) [Read more](#)

## Leibniz-HKI-Authors



Kerstin Kaerger

[Details](#)



**Oliver Kurzai**

[Details](#)



Grit Walther

[Details](#)

## Identifier

**doi:** 10.1016/j.mmcr.2019.07.008

**PMID:** 31428554