

Two-dimensional proteome reference maps for the human pathogenic filamentous fungus *Aspergillus fumigatus*.

Vödisch M, Albrecht D, Lessing F, Schmidt AD, Winkler R, Guthke R, Brakhage AA, Kniemeyer O (2009) Two-dimensional proteome reference maps for the human pathogenic filamentous fungus *Aspergillus fumigatus*. *Proteomics* 9(5), 1407-1415.

[Details](#)



Abstract

The filamentous fungus *Aspergillus fumigatus* has become the most important airborne fungal pathogen causing life-threatening infections in immunosuppressed patients. We established a 2-D reference map for *A. fumigatus*. Using MALDI-TOF-MS/MS, we identified 381 spots representing 334 proteins. Proteins involved in cellular metabolism, protein synthesis, transport processes and cell cycle were most abundant. Furthermore, we established a protocol for the isolation of mitochondria of *A. fumigatus* and developed a mitochondrial proteome reference map. 147 proteins represented by 234 spots were identified.

Involved units

[Molecular and Applied Microbiology Axel Brakhage](#) [Read more](#)

[Microbiome Dynamics Gianni Panagiotou](#) [Read more](#)

[International Leibniz Research School Peter F. Zipfel](#) [Read more](#)

Leibniz-HKI-Authors



Daniela Albrecht-Eckardt

[Details](#)



Axel A. Brakhage

[Details](#)



Reinhard Guthke

[Details](#)



Olaf Kniemeyer

[Details](#)



Franziska Lessing

[Details](#)



Martin Vödisch

[Details](#)

Topics

[Management of heterogeneous experimental data](#)

Identifier

doi: 10.1002/pmic.200800394

PMID: 19253289