

Microevolution of *Candida albicans* in macrophages restores filamentation in a nonfilamentous mutant.

Wartenberg A, Linde J, Martin R, Schreiner M, Horn F, Jacobsen ID, Jenull S, Wolf T, Kuchler K, Guthke R, Kurzai O, Forche A, d'Enfert C, Brunke S, Hube B (2014) Microevolution of *Candida albicans* in macrophages restores filamentation in a nonfilamentous mutant. *PLOS Genet* 10(12), e1004824.

[Details](#)



Abstract

Following antifungal treatment, *Candida albicans*, and other human pathogenic fungi can undergo microevolution, which leads to the emergence of drug resistance. However, the capacity for microevolutionary adaptation of fungi goes beyond the development of resistance against antifungals. Here we used an experimental microevolution approach to show that one of the central pathogenicity mechanisms of *C. albicans*, the yeast-to-hyphae transition, can be subject to experimental evolution. The *C. albicans* *cph1Δ/efg1Δ* mutant is nonfilamentous, as central signaling pathways linking environmental cues to hyphal formation are disrupted. We subjected this mutant to constant selection pressure in the hostile environment of the macrophage phagosome. In a comparatively short time-frame, the mutant evolved the ability to escape macrophages by filamentation. In addition, the evolved mutant exhibited hyper-virulence in a murine infection model and an altered cell wall composition compared to the *cph1Δ/efg1Δ* strain.

Moreover, the transcriptional regulation of hyphae-associated, and other pathogenicity-related genes became re-responsive to environmental cues in the evolved strain. We went on to identify the causative missense mutation via whole genome- and transcriptome-sequencing: a single nucleotide exchange took place within SSN3 that encodes a component of the Cdk8 module of the Mediator complex, which links transcription factors with the general transcription machinery. This mutation was responsible for the reconnection of the hyphal growth program with environmental signals in the evolved strain and was sufficient to bypass Efg1/Cph1-dependent filamentation. These data demonstrate that even central transcriptional networks can be remodeled very quickly under appropriate selection pressure.

Involved units

[Microbial Pathogenicity Mechanisms Bernhard Hube](#) [Read more](#)

[Fungal Septomics Oliver Kurzai](#) [Read more](#)

[Microbial Immunology Ilse Jacobsen](#) [Read more](#)

[Microbiome Dynamics Gianni Panagiotou](#) [Read more](#)

Leibniz-HKI-Authors



Sascha Brunke

[Details](#)



Reinhard Guthke

[Details](#)



Fabian Horn

[Details](#)



Bernhard Hube

[Details](#)



Ilse Denise Jacobsen

[Details](#)



Oliver Kurzai

[Details](#)



Jörg Linde

[Details](#)



Ronny Martin

[Details](#)



Maria Schreiner

[Details](#)



Anja Wartenberg

[Details](#)



Thomas Wolf

[Details](#)

Topics

[Evolution & adaptation in pathogenicity](#)

Awards

Recommended by Faculty of 1000 "[?][?][?]"; Medac Research Award 2015

Identifier

doi: 10.1371/journal.pgen.1004824

