

***Candida albicans*-epithelial interactions and induction of mucosal innate immunity.**

Naglik JR, König A, Hube B, Gaffen SL (2017) *Candida albicans*-epithelial interactions and induction of mucosal innate immunity. *Curr Opin Microbiol* 40, 104-112. (Review)

[Details](#)



Abstract

Candida albicans is a human fungal pathogen that causes millions of mucosal and life-threatening infections annually. *C. albicans* initially interacts with epithelial cells, resulting in fungal recognition and the formation of hyphae. Hypha formation is critical for host cell damage and immune activation, which are both driven by the secretion of Candidalysin, a recently discovered peptide toxin. Epithelial activation leads to the production of inflammatory mediators that recruit innate immune cells including neutrophils, macrophages and innate Type 17 cells, which together work with epithelial cells to clear the fungal infection. This review will focus on the recent discoveries that have advanced our understanding of *C. albicans*-epithelial interactions and the induction of mucosal innate immunity.

Involved units

[Microbial Pathogenicity Mechanisms](#) [Bernhard Hube](#) [Read more](#)

Leibniz-HKI-Authors



Bernhard Hube

[Details](#)



Annika König

[Details](#)

Identifier

doi: 10.1016/j.mib.2017.10.030

PMID: 29156234