

The *Candida albicans* exotoxin Candidalysin promotes alcohol-associated liver disease.

Chu H, Duan Y, Lang S, Jiang L, Wang Y, Llorente C, Liu J, Mogavero S, Bosques-Padilla F, Abalde JG, Vargas V, Tu XM, Yang L, Hou X, Hube B, Stärkel P, Schnabl B (2019) The *Candida albicans* exotoxin Candidalysin promotes alcohol-associated liver disease. *J Hepatol* 72(3), 391-400.

[Details](#)



Abstract

Alcohol-associated liver disease is a leading indication for liver transplantation and leading cause of mortality. Alterations of the gut microbiota contribute to pathogenesis of alcohol-associated liver disease. Patients with alcohol-associated liver disease have increased proportions of *Candida* spp. in the fecal mycobiome. However, little is known about the effect of intestinal *Candida* on alcohol-associated liver disease. Here we evaluated the contributions of *Candida albicans* and its exotoxin Candidalysin on alcoholic liver disease.

Involved units

[Microbial Pathogenicity Mechanisms Bernhard Hube](#) [Read more](#)

Leibniz-HKI-Authors



Bernhard Hube

[Details](#)



Selene Mogavero

[Details](#)

Topics

[Damage to the host](#)

[Evolution & adaptation in pathogenicity](#)

[Nutrient acquisition in infections](#)

Identifier

doi: 10.1016/j.jhep.2019.09.029

PMID: 31606552