

HapX-mediated adaption to iron starvation is crucial for virulence of *Aspergillus fumigatus*.

Schrettl M, Beckmann N, Varga J, Heinekamp T, Jacobsen ID, Jöchl C, Moussa TA, Wang S, Gsaller F, Blatzer M, Werner ER, Niermann WC, Brakhage AA, Haas H (2010) HapX-mediated adaption to iron starvation is crucial for virulence of *Aspergillus fumigatus*. *PLOS Pathog* 6(9), e1001124-e1001124.

[Details](#)



Abstract

Iron is essential for a wide range of cellular processes. Here we show that the bZIP-type regulator HapX is indispensable for the transcriptional remodeling required for adaption to iron starvation in the opportunistic fungal pathogen *Aspergillus fumigatus*. HapX represses iron-dependent and mitochondrial-localized activities including respiration, TCA cycle, amino acid metabolism, iron-sulfur-cluster and heme biosynthesis. In agreement with the impact on mitochondrial metabolism, HapX-deficiency decreases resistance to tetracycline and increases mitochondrial DNA content. Pathways positively affected by HapX include production of the ribotoxin AspF1 and siderophores, which are known virulence determinants. Iron starvation causes a massive remodeling of the amino acid pool and HapX is essential for the coordination of the production of siderophores and their precursor ornithine. Consistent with HapX-function being limited to iron depleted conditions and *A. fumigatus* facing iron starvation in the host, HapX-deficiency causes significant attenuation of virulence in a murine model of aspergillosis. Taken together, this study demonstrates that HapX-

dependent adaption to conditions of iron starvation is crucial for virulence of *A. fumigatus*.

Involved units

[Molecular and Applied Microbiology Axel Brakhage](#) [Read more](#)

[Microbial Immunology Ilse Jacobsen](#) [Read more](#)

Leibniz-HKI-Authors



Axel A. Brakhage

[Details](#)



Thorsten Heinekamp

[Details](#)



Ilse Denise Jacobsen

[Details](#)

Identifier

doi: 10.1371/journal.ppat.1001124

PMID: 20941352