

Phytotoxin production in *Aspergillus terreus* is regulated by independent environmental signals.

Gressler M, Meyer F, Heine D, Hortschansky P, Hertweck C, Brock M (2015) Phytotoxin production in *Aspergillus terreus* is regulated by independent environmental signals. *eLife* 4, e07861.

[Details](#)



Abstract

Secondary metabolites have a great potential as pharmaceuticals, but there are only few examples where regulation of gene cluster expression has been correlated with ecological and physiological relevance for the producer. Here, signals, mediators and biological effects of terrein production were studied in the fungus *Aspergillus terreus* to elucidate the contribution of terrein to ecological competition. Terrein causes fruit surface lesions and inhibits plant seed germination. Additionally, terrein is moderately antifungal and reduces ferric iron, thereby supporting growth of *A. terreus* under iron starvation. In accordance, the lack of nitrogen or iron or elevated methionine levels induced terrein production and was dependent on either the nitrogen response regulators AreA and AtfA or the iron response regulator HapX. Independent signal transduction allows complex sensing of the environment and, combined with its broad spectrum of biological activities, terrein

provides a prominent example for adapted secondary metabolite production in response to environmental competition.

Involved units

[Biomolecular Chemistry](#) [Christian Hertweck](#) [Read more](#)

[Molecular and Applied Microbiology](#) [Axel Brakhage](#) [Read more](#)

[Microbial Biochemistry and Physiology](#) [Read more](#)

Leibniz-HKI-Authors



Matthias Brock

[Details](#)



Markus Greßler

[Details](#)



Daniel Heine

[Details](#)



Christian Hertweck

[Details](#)



Peter Hortschansky

[Details](#)



Florian Meyer

[Details](#)

Topics

[Iron uptake and transport](#)

[Novel bacterial siderophores](#)

[Secondary metabolites from *A. terreus*](#)

Identifier

doi: 10.7554/eLife.07861

PMID: 26173180