

Reconstitution of iterative thioamidation in closthioamide biosynthesis reveals a novel nonribosomal peptide backbone-tailoring strategy.

Dunbar KL, Dell M, Molloy EM, Kloss F, Hertweck C (2019) Reconstitution of iterative thioamidation in closthioamide biosynthesis reveals a novel nonribosomal peptide backbone-tailoring strategy. *Angew Chem Int Ed* 58(37), 13014-13018.

[Details](#)



Abstract

Thioamide-containing nonribosomal peptides (NRPs) are exceedingly rare. Recently the biosynthetic gene cluster for the thioamidated NRP antibiotic closthioamide (CTA) was reported; however, the enzyme responsible for and timing of thioamide formation remained enigmatic. Here, we use genome editing, biochemical assays and mutational studies to demonstrate that a Fe-S cluster-containing member of the adenine nucleotide α -hydrolase protein superfamily (CtaC) is responsible for sulfur incorporation during CTA biosynthesis. However, unlike all previously characterized members, CtaC functions in a thiotemplated manner. The reconstitution of this unusual ATP-dependent sulfur transferase facilitates a revision of the CTA biosynthetic pathway and provides the first example of a NRP thioamide synthetase. Finally, using CtaC as a

bioinformatic handle, we provide promising clues that gene clusters encoding thioamidated NRPs are more widespread than previously thought.

Involved units

[Biomolecular Chemistry](#) [Christian Hertweck](#) [Read more](#)

[Transfer Group Anti-infectives](#) [Axel Brakhage](#) [Read more](#)

Leibniz-HKI-Authors



Maria Dell

[Details](#)



Kyle Dunbar

[Details](#)



Christian Hertweck

[Details](#)



Florian Kloß

[Details](#)



Evelyn Molloy

[Details](#)

Identifier

doi: 10.1002/anie.201905992

PMID: 31276268